Conditional Probability, Independence, Bayes Theorem 18.05 Spring 2014 Jeremy Orloff and Jonathan Bloom

Illustration of the Monty Hall problem removed due to copyright restrictions.

Sample Space Confusions

- 1. Sample space = set of all possible outcomes of an experiment.
- 2. The size of the set is not the sample space.
- 3. Outcomes can be sequences of numbers.

Examples.

1. Roll 5 dice: Ω = set of all sequences of 5 numbers between 1 and 6, e.g. $(1, 2, 1, 3, 1, 5) \in \Omega$. The size $|\Omega| = 6^5$ is not a set.

2.
$$\Omega = \text{set of all sequences of 10 birthdays,}$$


e.g. $(111, 231, 3, 44, 55, 129, 345, 14, 24, 14) \in \Omega$.
 $|\Omega| = 365^{10}$


3. *n* some number, $\Omega = \text{set of all sequences of } n$ birthdays. $|\Omega| = 365^n$. Don't forget that after class we post the slides including solutions to all the questions.

Conditional Probability

'the probability of A given B'.

$$P(A|B) = rac{P(A \cap B)}{P(B)}$$
, provided $P(B) \neq 0$.





Conditional probability: Abstractly and for coin example

Concept Question

Toss a coin 4 times. Let

- A = 'at least three heads'
- B = 'first toss is tails'.
- 1. What is P(A|B)?
- a) 1/16 b) 1/8 c) 1/4 d) 1/5

2. What is P(B|A)? a) 1/16 b) 1/8 c) 1/4 d) 1/5

Table Question

"Steve is very shy and withdrawn, invariably helpful, but with little interest in people, or in the world of reality. A meek and tidy soul, he has a need for order and structure and a passion for detail."*


What is the probability that Steve is a librarian? What is the probability that Steve is a farmer?

*From *Judgment under uncertainty: heuristics and biases* by Tversky and Kahneman.

Multiplication Rule, Law of Total Probability MR: $P(A \cap B) = P(A|B) \cdot P(B)$.

LoTP: If B_1 , B_2 , B_3 partition Ω then

 $P(A) = P(A \cap B_1) + P(A \cap B_2) + P(A \cap B_3)$ = $P(A|B_1)P(B_1) + P(A|B_2)P(B_2) + P(A|B_3)P(B_3)$




Trees


- Organize computations
- Compute total probability
- Compute Bayes formula

Example. : Game: 5 red and 2 green balls in an urn. A random ball is selected and replaced by a ball of the other color.

1. What is the probability the second ball is red?


2. What is the probability the first ball was red given the second ball was red?






1. The probability x represents


a) $P(A_1)$ b) $P(A_1|B_2)$ c) $P(B_2|A_1)$ d) $P(C_1|B_2 \cap A_1)$.



- 2. The probability y represents
- a) $P(B_2)$ b) $P(A_1|B_2)$ c) $P(B_2|A_1)$ d) $P(C_1|B_2 \cap A_1)$.



- 3. The probability z represents
- a) $P(C_1)$ b) $P(B_2|C_1)$ c) $P(C_1|B_2)$ d) $P(C_1|B_2 \cap A_1)$.



- 4. The circled node represents the event
- a) C_1 b) $B_2 \cap C_1$ c) $A_1 \cap B_2 \cap C_1$ d) $C_1|B_2 \cap A_1$.

Concept question: Monty Hall

Let's Make a Deal

- 1. One door hides a car, two hide goats.
- 2. Contestant chooses door.
- 3. Monty (who knows where the car is) opens a different door with a goat.
- 4. Contestant can switch doors or keep her original choice. What is the best strategy for winning a car?
- a) Switch b) Don't switch c) It doesn't matter

Illustration of the Monty Hall problem removed due to copyright restrictions.

- Organize the Monty Hall problem into a tree and compute the probability of winning if you always switch.
- Hint first break the game into a sequence of actions.

Independence

Events A and B are independent if the probability that one occurred is not affected by knowledge that the other occurred.

A and B are independent $\Leftrightarrow P(A \cap B) = P(A)P(B)$ $\Leftrightarrow P(A|B) = P(A)$ $\Leftrightarrow P(B|A) = P(B)$

Concept Question: Independence

Roll two dice.

- A = 'first die is 3'
- B = `sum is 6'

A and B are independent a) True b) False Also called Bayes Rule and Bayes Formula.

Allows you to find P(A|B) from P(B|A), i.e. to 'invert' conditional probabilities.

$$P(A|B) = rac{P(B|A) \cdot P(A)}{P(B)}$$

Often compute the denominator P(B) using the law of total probability.

Board Question: Evil Squirrels

Of the one million squirrels on MIT's campus most are good-natured. But one hundred of them are pure evil! An enterprising student in Course 6 develops an "Evil Squirrel Alarm" which she offers to sell to MIT for a passing grade. MIT decides to test the reliability of the alarm by conducting trials.

Photograph of squirrel removed due to copyright restrictions.

Evil Squirrels Continued

- When presented with an evil squirrel, the alarm goes off 99% of the time.
- When presented with a good-natured squirrel, the alarm goes off 1% of the time.
- a) If a squirrel sets off the alarm, what is the probability that it is evil?
- b) Should MIT co-opt the patent rights and employ the system?

One solution

| | Evil | Nice | |
|----------|------|--------|---------|
| Alarm | 99 | 9999 | 10098 |
| No alarm | 1 | 989901 | 989902 |
| | 100 | 999900 | 1000000 |

Summary:

Probability a random test is correct $= \frac{99+989901}{1000000} = .99$

Probability a positive test is correct $= \frac{99}{10098} \approx .01$

These probabilities are not the same!

Washington Post, hot off the press

Annual physical exam is probably unnecessary if you're generally healthy

For patients, the negatives include time away from work and possibly unnecessary tests. "Getting a simple urinalysis could lead to a false positive, which could trigger a cascade of even more tests, only to discover in the end that you had nothing wrong with you." Mehrotra says.

http://www.washingtonpost.com/national/health-science/ annual-physical-exam-is-probably-unnecessary-if-youregenerally-healthy/2013/02/08/2c1e326a-5f2b-11e2-a389ee565c81c565_story.html

18.05 Introduction to Probability and Statistics Spring 2014

For information about citing these materials or our Terms of Use, visit: http://ocw.mit.edu/terms.