

18.905 Problem Set 6

Due Wednesday, October 18 in class

1. Use the universal coefficient theorem to compute $H_*(L(p, q); \mathbb{Z}/m)$ for all m , where $L(p, q)$ are the lens spaces defined in class.
2. Hatcher, Exercise 1 on page 267.
3. Given an arbitrary finitely generated abelian group M , compute $\text{Tor}_k(\mathbb{Q}/\mathbb{Z}, M)$ for all $k \geq 0$. (Bonus marks for doing an arbitrary abelian group.)
4. The n -dimensional chains C_n form a functor from spaces to abelian groups. Suppose F is another functor from spaces to abelian groups. Show that any natural transformation Θ from C_n to F must be as follows: For any space X , the map $\Theta_X : C_n(X) \rightarrow F(X)$ is given by

$$\Theta_X \left(\sum m_\sigma \sigma \right) = \sum m_\sigma F(\sigma)(\Theta_{\Delta^n} \Delta^n).$$

(Here we recall that the chain $\Delta^n \in C_n(\Delta^n)$ is represented by the identity map from Δ^n to itself.) Conversely, given an element $S \in F(\Delta^n)$, show that we can define a natural transformation Θ from C_n to F by

$$\Theta_X \left(\sum m_\sigma \sigma \right) = \sum m_\sigma F(\sigma)(S).$$